

Dyсен *Forte Tablet / Suspension*

(Secnidazole)

COMPOSITION:**Dyсен Forte Tablet:**

Each film coated tablet contains:
Secnidazole ... 1000mg.
NQ Specs.

Dyсен 750mg Suspension:

When reconstituted as directed:
Each 30ml contains:
Secnidazole ... 750mg
as granules for suspension.
NQ Specs.

DESCRIPTION:

Secnidazole is a synthetic derivative of the nitroimidazole series.

PHARMACOKINETICS:

The maximum serum level is obtained after 3 hours following oral administration of 2g Secnidazole. The plasma elimination half-life is about 25 hours. The majority of **Secnidazole** is eliminated via urine (50% of the administered dose is excreted within 12 hours). Secnidazole crosses the placental barrier and can be found in maternal milk.

INDICATIONS:

Dyсен is indicated for the treatment of Intestinal amoebiasis, Hepatic amoebiasis, Giardiasis, Urethritis and Vaginitis due to *Trichomonas vaginalis*.

CONTRA-INDICATIONS:

The product should not be administered during first trimester of pregnancy or during lactation and hypersensitivity to imidazole derivatives.

DOSAGE AND ADMINISTRATION:

The dose should preferably be taken during a meal.

Acute Intestinal Amoebiasis and Giardiasis:

Adults: Single dose treatment of two **Secnidazole** Tablets (2g).
Children over 2 years: Single dose of 30mg/kg/day, with a maximum of 2000mg.

Hepatic Amoebiasis:

Adult: One and a half (1½) tablets of 1000mg (1.5g) as a single dose for 5-7 days.

Children: 30mg/kg body weight once a day or in divided doses before meals for 5-7 days.

In suppurative hepatic amoebiasis, Secnidazole therapy should be complemented by abscess drainage.

Urogenital Trichomoniasis and Bacterial vaginosis:

Single dose treatment of two **Secnidazole** Tablets (2g). The male partner should also be treated with the same dose.

DIRECTIONS FOR PREPARATION OF DYSEN SUSPENSION:

Loosen the powder by shaking the bottle. To prepare 30ml oral suspension, use provided measuring cup of 10ml. First add a cup of 10ml previously boiled and cool water, close cap and shake vigorously. After this once again add another cup of 10ml of previously boiled and cool water, recap the bottle and again shake bottle vigorously for final preparation of suspension for use. Take immediately after preparation in a single dose.

PRECAUTIONS:

Avoid alcohol during treatment. Do not associate with disulfiram. Avoid during first trimester of pregnancy. Do not administer to subjects with history of blood dyscrasia. This product is not recommended in children below 2 years of age.

DRUG AND OTHER INTERACTIONS:

- Contra-indicated combination therapy
- With Disulfiram: delirious episodes, confusion.
 - With Alcohol: hotness, redness, vomiting, tachycardia.

COMBINATIONS SUBJECT TO PRECAUTIONS:

Oral anticoagulant (described for warfarin): potentiation of the oral anticoagulant and increased risk of haemorrhage by lowering of its hepatic catabolism. More frequent determination of the prothrombin time and monitoring of the International normalized ratio (INR). Adjustment of the oral anticoagulant dose during Secnidazole treatment, and 8 days after its withdrawal.

Rev. 05-2015

SIDE EFFECTS:

The side effects of **Secnidazole** are same as that of other nitroimidazole derivatives.

- Most frequently: digestive disorders with abdominal pain, change in taste (metallic), glossitis, stomatitis.
- Urticaria.
- Moderate leukopenia, reversible upon withdrawal of treatment, most rarely, dizziness, lack of co-ordination and ataxia, paraesthesia, sensitive and motor polyneuritis with Secnidazole. Rare cases of digestive disorders have been reported (nausea, vomiting, gastric pain).

OVERDOSAGE:

Symptoms: Expected symptoms are drowsiness, dizziness headache, ataxia, skin rashes, pruritus, transient epileptiform, seizure.

TREATMENT:

There is no specific treatment for gross overdosage. The treatment is symptomatic and supportive.

INSTRUCTIONS:

Store below 30°C. Protect from heat, light and moisture. Keep out of the reach of children. **Shake well before use.**

PRESENTATION:

Dysen Forte Tablets: Blister Pack of 1x2 tablets.

Dysen 750mg Suspension: Pack of 1 bottle contains granules for suspension.

ڈائیسین فورٹ گولیاں / (سیکنڈازول)

سیکنڈازول ایک جدید تحقیقی دوا ہے جو کہ مندرجہ ذیل علامات میں مرینس کے لئے تجویز کی جاتی ہے۔

امیبیاس

جیبارڈیاس

بیکٹیریل ویجیٹوس

ٹراکیمونی ایس

ڈائیسین سسپنشن تیار کرنے کا طریقہ: بوتل کو گھسی طرح بلائیں تاکہ تمام پاؤڈر نرم ہو جائے۔ ۳۰ ملی لیٹر سسپنشن بنانے کے لئے ڈبے میں موجود کپ سے ۱۰ ملی لیٹر لیا ہوا ٹھنڈا پانی بھر کر بوتل میں ڈالیں اور بوتل کو گھسی طرح بلائیں۔ اس کے بعد مزید ایک کپ پانی (۱۰ ملی لیٹر) بوتل میں ڈال کر بوتل کو گھسی طرح بلائیں تاکہ تمام پاؤڈر پانی میں حل ہو جائے۔ اس طرح تیار کی ہوئی دوا کو ایک ہی خوراک میں فوراً استعمال کریں۔

خواص: سیکنڈازول کی ایک خوراک مندرجہ بالا علامات کے لئے مکمل علاج ہے:

سیکنڈازول کی ایک خوراک علاج مرض کی تمام علامات کو مکمل طور پر ختم کر کے اس کے دوبارہ ہونے کے امکانات کا خاتمہ کرتا ہے۔

سیکنڈازول کی ایک خوراک کی بدولت یورٹرائٹس اور ویجیٹائٹس کی علامات میں دونوں ساتھیوں (Partners) کے علاج میں معاون ثابت ہوتا ہے۔

سیکنڈازول سسپنشن بچوں میں علاج کی وہ سہولت فراہم کرتا ہے جو کہ کثیر مقدار خوراک والی ادویات سے ممکن نہیں۔

ہدایات:

۳۰ ڈگری سینٹی گریڈ سے کم درجہ حرارت پر رکھیں۔ گرمی، روشنی اور نمی سے بچائیں۔ بچوں کی پہنچ سے دور رکھیں۔

استعمال سے پہلے بوتل کو گھسی طرح بلائیں۔



Manufactured by:
NABIQASIM INDUSTRIES (PVT) LTD.
17/24, Korangi Industrial Area,
Karachi-Pakistan.

Rev: 05-2015